

cordially invites you to attend a program entitled

EXPAREL® (bupivacaine liposome injectable suspension): An Opioid-Free Option for Postsurgical Pain

Presented by

James Mueller, MD
Staff Anesthesiologist
Medical City Hospital
Dallas, TX

Tuesday, May 8, 2018
6:30 PM

Olive & June
3411 Glenview Ave.
Austin, TX 78703
(512) 467-9898

Host: Carrie Crutchfield
(415) 806-6335; Carrie.Crutchfield@pacira.com

Please RSVP on or before Wednesday, May 2, 2018

Register online at www.PaciraPortal.com/Registration using Event ID 3780
or by completing the registration information below and sending via fax to (973) 577-8637
or via e-mail to crsreservation@pvaluecomm.com

If you have any questions about this program, please call p-value communications at (866) 782-5830

IMPORTANT: All fields are required. Please bring these numbers with you to the meeting.

Last Name _____ First Name _____
NPI Number _____ State of Licensure _____ State License # _____
Affiliated Institution _____
Phone _____ E-mail _____
Please indicate your credentials: MD DO DPM NP PA CRNA PharmD RN Other _____

Please see indication and Important Safety Information on the following page.

Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Adverse reactions reported with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation.

If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine.

EXPAREL is not recommended to be used in the following patient population: patients <18 years old and/or pregnant patients.

Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

[[PP-EX-US-3525 4/7/18 3:28 pm]]